

Morrison Institute / Cronkite News Poll Methodology and Results for Opioid Survey

The Morrison Institute-Cronkite News Poll surveyed 800 randomly selected Arizona adults regarding their opinions and behaviors surrounding opioid-based painkillers. The sample is representative of Arizona's adult population. Interviews were conducted March 11-18, 2017 and were about 11 minutes in length. The sample was quota-selected from 18 states based on age, gender and race to match the demographic characteristics of Arizona based on the latest Census data. The sampling frame included both landline- and cell-based telephones and interviews were conducted in Spanish as needed. The margin of error is not greater than 3.5 percentage points for the full sample.

Tab_agecat

Percent answers to 'Age' by 'All Respondents'

	Age 18-35		Age 36-64		Age 65+	
	Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 800)	33.0	(± 3.3)	47.0	(± 3.5)	20.0	(± 2.8)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_Q1

Percent answers to 'Just to be clear, when we ask about prescription painkillers in this poll, we mean strong ones, sometimes called opioids, such as Percocet, OxyContin or Vicodin. How easy or difficult do you think it is for people to get access to prescription painkillers?' by 'All Respondents'												
			Very easy		Somewhat easy		Somewhat difficult		Very difficult		Don't Know	
			Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 800)			29.6	(± 3.2)	28.6	(± 3.2)	15.4	(± 2.6)	8.4	(± 2.0)	18.0	(± 2.7)
Notes: 95 percent margin of error (MOE) in parentheses.												
Percent answers to 'Just to be clear, when we ask about prescription painkillers in this poll, we mean strong ones, sometimes called opioids, such as Percocet, OxyContin or Vicodin. How easy or difficult do you think it is for people to get access to prescription painkillers?' by 'Age'												
			Very easy		Somewhat easy		Somewhat difficult		Very difficult		Don't Know	
			Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE
Age 18-35 (N = 264)			29.9	(± 5.6)	36.7	(± 5.9)	17.0	(± 4.6)	4.9	(± 2.7)	11.4	(± 3.9)
Age 36-64 (N = 376)			30.3	(± 4.7)	25.8	(± 4.5)	15.2	(± 3.7)	10.4	(± 3.1)	18.4	(± 4.0)
Age 65+ (N = 160)			27.5	(± 7.1)	21.9	(± 6.5)	13.1	(± 5.3)	9.4	(± 4.6)	28.1	(± 7.1)

Notes: 95 percent margin of error (MOE) in parentheses.												
Percent answers to 'Just to be clear, when we ask about prescription painkillers in this poll, we mean strong ones, sometimes called opioids, such as Percocet, OxyContin or Vicodin. How easy or difficult do you think it is for people to get access to prescription painkillers?' by 'Gender'												
			Very easy		Somewhat easy		Somewhat difficult		Very difficult		Don't Know	
			Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE
Female (N = 404)			29.2	(± 4.5)	29.7	(± 4.5)	14.4	(± 3.5)	7.7	(± 2.7)	19.1	(± 3.9)
Male (N = 394)			30.2	(± 4.6)	27.4	(± 4.5)	16.5	(± 3.7)	8.9	(± 2.9)	17.0	(± 3.8)
Notes: 95 percent margin of error (MOE) in parentheses.												
Percent answers to 'Just to be clear, when we ask about prescription painkillers in this poll, we mean strong ones, sometimes called opioids, such as Percocet, OxyContin or Vicodin. How easy or difficult do you think it is for people to get access to prescription painkillers?' by 'Race'												
			Very easy		Somewhat easy		Somewhat difficult		Very difficult		Don't Know	
			Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE
Black (N = 28)			35.7	(±18.1)	35.7	(±18.1)	7.1	(± 9.7)	0.0	(± 0.0)	21.4	(±15.5)
Hispanic (N = 208)			27.9	(± 6.2)	26.4	(± 6.1)	15.9	(± 5.1)	10.6	(± 4.3)	19.2	(± 5.5)
Other race (N = 76)			31.6	(±10.7)	27.6	(±10.3)	15.8	(± 8.4)	7.9	(± 6.2)	17.1	(± 8.6)
White (N = 488)			29.7	(± 4.1)	29.3	(± 4.1)	15.6	(± 3.3)	8.0	(± 2.5)	17.4	(± 3.4)
Notes: 95 percent margin of error (MOE) in parentheses.												

Tab_Q2

Percent answers to 'To the best of your knowledge, are heroin and prescription painkillers similar or different types of drugs in terms of how they affect the drug user's body?' by 'All Respondents'								
			Similar		Different		Don't Know	
			Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 799)			47.3	(± 3.5)	22.7	(± 3.0)	30.0	(± 3.2)
Notes: 95 percent margin of error (MOE) in parentheses.								
Percent answers to 'To the best of your knowledge, are heroin and prescription painkillers similar or different types of drugs in terms of how they affect the drug user's body?' by 'Age'								
			Similar		Different		Don't Know	

		Percent	MOE	Percent	MOE	Percent	MOE
Age 18-35 (N = 264)		57.2	(± 6.1)	22.0	(± 5.1)	20.8	(± 5.0)
Age 36-64 (N = 375)		46.1	(± 5.2)	23.2	(± 4.4)	30.7	(± 4.8)
Age 65+ (N = 160)		33.8	(± 7.5)	22.5	(± 6.6)	43.8	(± 7.8)
Notes: 95 percent margin of error (MOE) in parentheses.							
Percent answers to 'To the best of your knowledge, are heroin and prescription painkillers similar or different types of drugs in terms of how they affect the drug user's body?' by 'Gender'							
		Similar		Different		Don't Know	
		Percent	MOE	Percent	MOE	Percent	MOE
Female (N = 403)		47.6	(± 5.0)	20.8	(± 4.0)	31.5	(± 4.6)
Male (N = 394)		47.0	(± 5.0)	24.6	(± 4.3)	28.4	(± 4.5)
Notes: 95 percent margin of error (MOE) in parentheses.							
Percent answers to 'To the best of your knowledge, are heroin and prescription painkillers similar or different types of drugs in terms of how they affect the drug user's body?' by 'Race'							
		Similar		Different		Don't Know	
		Percent	MOE	Percent	MOE	Percent	MOE
Black (N = 28)		53.6	(±18.9)	14.3	(±13.2)	32.1	(±17.7)
Hispanic (N = 207)		45.9	(± 6.9)	22.7	(± 5.8)	31.4	(± 6.5)
Other race (N = 76)		46.1	(±11.4)	23.7	(± 9.8)	30.3	(±10.5)
White (N = 488)		47.7	(± 4.5)	23.0	(± 3.8)	29.3	(± 4.1)
Notes: 95 percent margin of error (MOE) in parentheses.							

Tab_Q3

Percent answers to 'Which do you think is riskier to use, prescription painkillers or heroin, or do you think they are about equally risky?' by 'All Respondents'									
		Prescription Painkillers		Heroin		About equal		Don't Know	
		Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 800)		6.5	(± 1.7)	32.6	(± 3.3)	54.2	(± 3.5)	6.6	(± 1.8)
Notes: 95 percent margin of error (MOE) in parentheses.									
Percent answers to 'Which do you think is riskier to use, prescription painkillers or heroin, or do you think they are about equally risky?' by 'Age'									

			Prescription Painkillers		Heroin		About equal		Don't Know	
			Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE
Age 18-35 (N = 264)			3.8	(± 2.4)	35.2	(± 5.9)	58.3	(± 6.1)	2.7	(± 2.0)
Age 36-64 (N = 376)			9.6	(± 3.0)	31.4	(± 4.8)	53.5	(± 5.1)	5.6	(± 2.4)
Age 65+ (N = 160)			3.8	(± 3.0)	31.2	(± 7.3)	49.4	(± 7.9)	15.6	(± 5.7)
Notes: 95 percent margin of error (MOE) in parentheses.										
Percent answers to 'Which do you think is riskier to use, prescription painkillers or heroin, or do you think they are about equally risky?' by 'Gender'										
			Prescription Painkillers		Heroin		About equal		Don't Know	
			Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE
Female (N = 404)			5.2	(± 2.2)	29.0	(± 4.5)	59.2	(± 4.9)	6.7	(± 2.5)
Male (N = 394)			7.9	(± 2.7)	36.3	(± 4.8)	49.2	(± 5.0)	6.6	(± 2.5)
Notes: 95 percent margin of error (MOE) in parentheses.										
Percent answers to 'Which do you think is riskier to use, prescription painkillers or heroin, or do you think they are about equally risky?' by 'Race'										
			Prescription Painkillers		Heroin		About equal		Don't Know	
			Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE
Black (N = 28)			7.1	(± 9.7)	28.6	(±17.1)	57.1	(±18.7)	7.1	(± 9.7)
Hispanic (N = 208)			12.5	(± 4.6)	26.4	(± 6.1)	56.2	(± 6.9)	4.8	(± 3.0)
Other race (N = 76)			3.9	(± 4.5)	36.8	(±11.1)	48.7	(±11.5)	10.5	(± 7.0)
White (N = 488)			4.3	(± 1.8)	34.8	(± 4.3)	54.1	(± 4.5)	6.8	(± 2.3)
Notes: 95 percent margin of error (MOE) in parentheses.										

Tab_Q4

Percent answers to 'Do you think prescription painkiller abuse makes a person more likely or less likely to use heroin or other illegal drugs, or do you think it doesn't make much of a difference?' by 'All Respondents'										
			More Likely		Less Likely		No Difference		Don't Know	
			Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 798)			58.1	(± 3.5)	3.1	(± 1.2)	27.7	(± 3.2)	11.0	(± 2.2)
Notes: 95 percent margin of error (MOE) in parentheses.										

Percent answers to 'Do you think prescription painkiller abuse makes a person more likely or less likely to use heroin or other illegal drugs, or do you think it doesn't make much of a difference?' by 'Age'										
			More Likely		Less Likely		No Difference		Don't Know	
			Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE
Age 18-35 (N = 264)			70.5	(± 5.6)	1.9	(± 1.7)	22.3	(± 5.1)	5.3	(± 2.8)
Age 36-64 (N = 374)			54.3	(± 5.2)	4.0	(± 2.0)	29.9	(± 4.7)	11.8	(± 3.3)
Age 65+ (N = 160)			46.9	(± 7.9)	3.1	(± 2.8)	31.2	(± 7.3)	18.8	(± 6.2)
Notes: 95 percent margin of error (MOE) in parentheses.										
Percent answers to 'Do you think prescription painkiller abuse makes a person more likely or less likely to use heroin or other illegal drugs, or do you think it doesn't make much of a difference?' by 'Gender'										
			More Likely		Less Likely		No Difference		Don't Know	
			Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE
Female (N = 404)			61.6	(± 4.8)	2.7	(± 1.6)	25.5	(± 4.3)	10.1	(± 3.0)
Male (N = 392)			54.3	(± 5.0)	3.6	(± 1.9)	30.1	(± 4.6)	12.0	(± 3.3)
Notes: 95 percent margin of error (MOE) in parentheses.										
Percent answers to 'Do you think prescription painkiller abuse makes a person more likely or less likely to use heroin or other illegal drugs, or do you think it doesn't make much of a difference?' by 'Race'										
			More Likely		Less Likely		No Difference		Don't Know	
			Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE
Black (N = 28)			60.7	(±18.5)	7.1	(± 9.7)	28.6	(±17.1)	3.6	(± 7.0)
Hispanic (N = 207)			57.5	(± 6.9)	3.4	(± 2.5)	27.1	(± 6.2)	12.1	(± 4.5)
Other race (N = 76)			60.5	(±11.2)	3.9	(± 4.5)	26.3	(±10.1)	9.2	(± 6.6)
White (N = 487)			57.9	(± 4.5)	2.7	(± 1.5)	28.1	(± 4.1)	11.3	(± 2.9)
Notes: 95 percent margin of error (MOE) in parentheses.										

Tab_Q5

Percent answers to 'In the past 12 months have you experienced ongoing pain for more than three months?' by 'All Respondents'										
			Yes		No		Don't Know			
			Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE

All Respondents (N = 799)	29.0	(± 3.2)	70.7	(± 3.2)	0.3	(± 0.4)
Notes: 95 percent margin of error (MOE) in parentheses.						
Percent answers to 'In the past 12 months have you experienced ongoing pain for more than three months?' by 'Age'						
	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Age 18-35 (N = 264)	15.2	(± 4.4)	84.8	(± 4.4)	0.0	(± 0.0)
Age 36-64 (N = 376)	39.4	(± 5.0)	60.4	(± 5.0)	0.3	(± 0.5)
Age 65+ (N = 159)	27.7	(± 7.1)	71.7	(± 7.1)	0.6	(± 1.3)
Notes: 95 percent margin of error (MOE) in parentheses.						
Percent answers to 'In the past 12 months have you experienced ongoing pain for more than three months?' by 'Gender'						
	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Female (N = 403)	29.5	(± 4.5)	70.5	(± 4.5)	0.0	(± 0.0)
Male (N = 394)	28.4	(± 4.5)	71.1	(± 4.6)	0.5	(± 0.7)
Notes: 95 percent margin of error (MOE) in parentheses.						
Percent answers to 'In the past 12 months have you experienced ongoing pain for more than three months?' by 'Race'						
	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Black (N = 28)	21.4	(±15.5)	78.6	(±15.5)	0.0	(± 0.0)
Hispanic (N = 208)	21.2	(± 5.7)	78.8	(± 5.7)	0.0	(± 0.0)
Other race (N = 76)	27.6	(±10.3)	72.4	(±10.3)	0.0	(± 0.0)
White (N = 487)	33.1	(± 4.3)	66.5	(± 4.3)	0.4	(± 0.6)
Notes: 95 percent margin of error (MOE) in parentheses.						

Tab_Q6

Percent answers to 'Are you in chronic pain?' by 'All Respondents'						
	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 232)	70.3	(± 6.0)	29.3	(± 6.0)	0.4	(± 0.9)

Notes: 95 percent margin of error (MOE) in parentheses.							
Percent answers to 'Are you in chronic pain?' by 'Age'							
		Yes		No		Don't Know	
		Percent	MOE	Percent	MOE	Percent	MOE
Age 18-35 (N = 40)		77.5	(±13.2)	22.5	(±13.2)	0.0	(± 0.0)
Age 36-64 (N = 148)		74.3	(± 7.2)	25.0	(± 7.1)	0.7	(± 1.3)
Age 65+ (N = 44)		50.0	(±15.1)	50.0	(±15.1)	0.0	(± 0.0)
Notes: 95 percent margin of error (MOE) in parentheses.							
Percent answers to 'Are you in chronic pain?' by 'Gender'							
		Yes		No		Don't Know	
		Percent	MOE	Percent	MOE	Percent	MOE
Female (N = 119)		70.6	(± 8.4)	29.4	(± 8.4)	0.0	(± 0.0)
Male (N = 112)		69.6	(± 8.7)	29.5	(± 8.6)	0.9	(± 1.8)
Notes: 95 percent margin of error (MOE) in parentheses.							
Percent answers to 'Are you in chronic pain?' by 'Race'							
		Yes		No		Don't Know	
		Percent	MOE	Percent	MOE	Percent	MOE
Black (N = 6)		100.0	(± 0.0)	0.0	(± 0.0)	0.0	(± 0.0)
Hispanic (N = 44)		65.9	(±14.3)	34.1	(±14.3)	0.0	(± 0.0)
Other race (N = 21)		66.7	(±20.6)	33.3	(±20.6)	0.0	(± 0.0)
White (N = 161)		70.8	(± 7.2)	28.6	(± 7.1)	0.6	(± 1.2)
Notes: 95 percent margin of error (MOE) in parentheses.							

Tab_Q7_cata_1

Percent answers to 'Over-the-counter pain relievers--In what ways did you try to relieve your pain?'						
by 'All Respondents'						
		Not Checked			Checked	
		Percent	MOE	Percent	MOE	

All Respondents (N = 232)	65.9	(± 6.2)	34.1	(± 6.2)
Notes: 95 percent margin of error (MOE) in parentheses.				
Percent answers to 'Over-the-counter pain relievers--In what ways did you try to relieve your pain?'				
by 'Age'				
		Not Checked		Checked
		Percent	MOE	Percent
				MOE
Age 18-35 (N = 40)	77.5	(±13.2)	22.5	(±13.2)
Age 36-64 (N = 148)	64.9	(± 7.9)	35.1	(± 7.9)
Age 65+ (N = 44)	59.1	(±14.9)	40.9	(±14.9)
Notes: 95 percent margin of error (MOE) in parentheses.				
Percent answers to 'Over-the-counter pain relievers--In what ways did you try to relieve your pain?'				
by 'Gender'				
		Not Checked		Checked
		Percent	MOE	Percent
				MOE
Female (N = 119)	64.7	(± 8.8)	35.3	(± 8.8)
Male (N = 112)	67.0	(± 8.9)	33.0	(± 8.9)
Notes: 95 percent margin of error (MOE) in parentheses.				
Percent answers to 'Over-the-counter pain relievers--In what ways did you try to relieve your pain?'				
by 'Race'				
		Not Checked		Checked
		Percent	MOE	Percent
				MOE
Black (N = 6)	83.3	(±30.5)	16.7	(±30.5)
Hispanic (N = 44)	63.6	(±14.5)	36.4	(±14.5)
Other race (N = 21)	71.4	(±19.8)	28.6	(±19.8)
White (N = 161)	65.2	(± 7.5)	34.8	(± 7.5)
Notes: 95 percent margin of error (MOE) in parentheses.				

Tab_Q7_cata_2

Percent answers to 'Prescription pain relievers--In what ways did you try to relieve your pain?'				
by 'All Respondents'				
		Not Checked		Checked

			Percent	MOE	Percent	MOE
All Respondents (N = 232)			64.2	(± 6.3)	35.8	(± 6.3)
Notes: 95 percent margin of error (MOE) in parentheses.						
Percent answers to 'Prescription pain relievers--In what ways did you try to relieve your pain?'						
by 'Age'						
			Not Checked		Checked	
			Percent	MOE	Percent	MOE
Age 18-35 (N = 40)			77.5	(±13.2)	22.5	(±13.2)
Age 36-64 (N = 148)			62.2	(± 8.0)	37.8	(± 8.0)
Age 65+ (N = 44)			59.1	(±14.9)	40.9	(±14.9)
Notes: 95 percent margin of error (MOE) in parentheses.						
Percent answers to 'Prescription pain relievers--In what ways did you try to relieve your pain?'						
by 'Gender'						
			Not Checked		Checked	
			Percent	MOE	Percent	MOE
Female (N = 119)			63.9	(± 8.8)	36.1	(± 8.8)
Male (N = 112)			65.2	(± 9.0)	34.8	(± 9.0)
Notes: 95 percent margin of error (MOE) in parentheses.						
Percent answers to 'Prescription pain relievers--In what ways did you try to relieve your pain?'						
by 'Race'						
			Not Checked		Checked	
			Percent	MOE	Percent	MOE
Black (N = 6)			33.3	(±38.6)	66.7	(±38.6)
Hispanic (N = 44)			72.7	(±13.5)	27.3	(±13.5)
Other race (N = 21)			71.4	(±19.8)	28.6	(±19.8)
White (N = 161)			62.1	(± 7.7)	37.9	(± 7.7)
Notes: 95 percent margin of error (MOE) in parentheses.						

Tab_Q7_cata_3

Percent answers to 'Chiropractic care--In what ways did you try to relieve your pain?'					
by 'All Respondents'					
		Not Checked		Checked	
		Percent	MOE	Percent	MOE
All Respondents (N = 232)		92.7	(± 3.4)	7.3	(± 3.4)
Notes: 95 percent margin of error (MOE) in parentheses.					
Percent answers to 'Chiropractic care--In what ways did you try to relieve your pain?'					
by 'Age'					
		Not Checked		Checked	
		Percent	MOE	Percent	MOE
Age 18-35 (N = 40)		85.0	(±11.3)	15.0	(±11.3)
Age 36-64 (N = 148)		94.6	(± 3.7)	5.4	(± 3.7)
Age 65+ (N = 44)		93.2	(± 7.6)	6.8	(± 7.6)
Notes: 95 percent margin of error (MOE) in parentheses.					
Percent answers to 'Chiropractic care--In what ways did you try to relieve your pain?'					
by 'Gender'					
		Not Checked		Checked	
		Percent	MOE	Percent	MOE
Female (N = 119)		91.6	(± 5.1)	8.4	(± 5.1)
Male (N = 112)		93.8	(± 4.6)	6.2	(± 4.6)
Notes: 95 percent margin of error (MOE) in parentheses.					
Percent answers to 'Chiropractic care--In what ways did you try to relieve your pain?'					
by 'Race'					
		Not Checked		Checked	

	Percent	MOE	Percent	MOE
Black (N = 6)	100.0	(± 0.0)	0.0	(± 0.0)
Hispanic (N = 44)	95.5	(± 6.3)	4.5	(± 6.3)
Other race (N = 21)	81.0	(±17.2)	19.0	(±17.2)
White (N = 161)	93.2	(± 4.0)	6.8	(± 4.0)
Notes: 95 percent margin of error (MOE) in parentheses.				

Tab_Q7_cata_4

Percent answers to 'Yoga/stretching--In what ways did you try to relieve your pain?'					
by 'All Respondents'					
	Not Checked			Checked	
	Percent	MOE	Percent	MOE	
All Respondents (N = 232)	93.1	(± 3.3)	6.9	(± 3.3)	
Notes: 95 percent margin of error (MOE) in parentheses.					
Percent answers to 'Yoga/stretching--In what ways did you try to relieve your pain?'					
by 'Age'					
	Not Checked			Checked	
	Percent	MOE	Percent	MOE	
Age 18-35 (N = 40)	85.0	(±11.3)	15.0	(±11.3)	
Age 36-64 (N = 148)	93.9	(± 3.9)	6.1	(± 3.9)	
Age 65+ (N = 44)	97.7	(± 4.5)	2.3	(± 4.5)	
Notes: 95 percent margin of error (MOE) in parentheses.					
Percent answers to 'Yoga/stretching--In what ways did you try to relieve your pain?'					
by 'Gender'					
	Not Checked			Checked	
	Percent	MOE	Percent	MOE	
Female (N = 119)	93.3	(± 4.6)	6.7	(± 4.6)	
Male (N = 112)	92.9	(± 4.9)	7.1	(± 4.9)	

Notes: 95 percent margin of error (MOE) in parentheses.					
Percent answers to 'Yoga/stretching--In what ways did you try to relieve your pain?'					
by 'Race'					
		Not Checked		Checked	
		Percent	MOE	Percent	MOE
Black (N = 6)		100.0	(± 0.0)	0.0	(± 0.0)
Hispanic (N = 44)		90.9	(± 8.7)	9.1	(± 8.7)
Other race (N = 21)		90.5	(±12.8)	9.5	(±12.8)
White (N = 161)		93.8	(± 3.8)	6.2	(± 3.8)
Notes: 95 percent margin of error (MOE) in parentheses.					

Tab_Q7_cata_5

Percent answers to 'Exercise--In what ways did you try to relieve your pain?'

by 'All Respondents'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
All Respondents (N = 232)	83.2	(± 4.9)	16.8	(± 4.9)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Exercise--In what ways did you try to relieve your pain?'

by 'Age'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Age 18-35 (N = 40)	82.5	(±12.0)	17.5	(±12.0)
Age 36-64 (N = 148)	82.4	(± 6.3)	17.6	(± 6.3)
Age 65+ (N = 44)	86.4	(±10.4)	13.6	(±10.4)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Exercise--In what ways did you try to relieve your pain?'

by 'Gender'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Female (N = 119)	78.2	(± 7.6)	21.8	(± 7.6)
Male (N = 112)	88.4	(± 6.1)	11.6	(± 6.1)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Exercise--In what ways did you try to relieve your pain?'

' by 'Race'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Black (N = 6)	100.0	(± 0.0)	0.0	(± 0.0)
Hispanic (N = 44)	70.5	(±13.8)	29.5	(±13.8)
Other race (N = 21)	90.5	(±12.8)	9.5	(±12.8)
White (N = 161)	85.1	(± 5.6)	14.9	(± 5.6)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_Q7_cata_6

Percent answers to 'Acupuncture--In what ways did you try to relieve your pain?'

' by 'All Respondents'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
All Respondents (N = 232)	98.3	(± 1.7)	1.7	(± 1.7)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Acupuncture--In what ways did you try to relieve your pain?'

' by 'Age'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Age 18-35 (N = 40)	97.5	(± 4.9)	2.5	(± 4.9)
Age 36-64 (N = 148)	98.6	(± 1.9)	1.4	(± 1.9)
Age 65+ (N = 44)	97.7	(± 4.5)	2.3	(± 4.5)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Acupuncture--In what ways did you try to relieve your pain?'

' by 'Gender'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Female (N = 119)	99.2	(± 1.7)	0.8	(± 1.7)
Male (N = 112)	97.3	(± 3.1)	2.7	(± 3.1)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Acupuncture--In what ways did you try to relieve your pain?'

by 'Race'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Black (N = 6)	100.0	(± 0.0)	0.0	(± 0.0)
Hispanic (N = 44)	97.7	(± 4.5)	2.3	(± 4.5)
Other race (N = 21)	100.0	(± 0.0)	0.0	(± 0.0)
White (N = 161)	98.1	(± 2.1)	1.9	(± 2.1)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_Q7_cata_7

Percent answers to 'Meditation--In what ways did you try to relieve your pain?'

by 'All Respondents'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
All Respondents (N = 232)	94.0	(± 3.1)	6.0	(± 3.1)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Meditation--In what ways did you try to relieve your pain?'

' by 'Age'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE

Age 18-35 (N = 40)	92.5	(± 8.3)	7.5	(± 8.3)
Age 36-64 (N = 148)	93.9	(± 3.9)	6.1	(± 3.9)
Age 65+ (N = 44)	95.5	(± 6.3)	4.5	(± 6.3)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Meditation--In what ways did you try to relieve your pain?

' by 'Gender'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Female (N = 119)	93.3	(± 4.6)	6.7	(± 4.6)
Male (N = 112)	94.6	(± 4.3)	5.4	(± 4.3)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Meditation--In what ways did you try to relieve your pain?

' by 'Race'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Black (N = 6)	100.0	(± 0.0)	0.0	(± 0.0)
Hispanic (N = 44)	93.2	(± 7.6)	6.8	(± 7.6)
Other race (N = 21)	100.0	(± 0.0)	0.0	(± 0.0)
White (N = 161)	93.2	(± 4.0)	6.8	(± 4.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q7_cata_8

Percent answers to 'Other ' by 'All Respondents'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
All Respondents (N = 232)	62.5	(± 6.4)	37.5	(± 6.4)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Other ' by 'Age'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Age 18-35 (N = 40)	62.5	(±15.3)	37.5	(±15.3)
Age 36-64 (N = 148)	60.1	(± 8.1)	39.9	(± 8.1)
Age 65+ (N = 44)	70.5	(±13.8)	29.5	(±13.8)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Other ' by 'Gender'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Female (N = 119)	60.5	(± 9.0)	39.5	(± 9.0)
Male (N = 112)	64.3	(± 9.1)	35.7	(± 9.1)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Other ' by 'Race'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Black (N = 6)	66.7	(±38.6)	33.3	(±38.6)
Hispanic (N = 44)	72.7	(±13.5)	27.3	(±13.5)
Other race (N = 21)	61.9	(±21.2)	38.1	(±21.2)
White (N = 161)	59.6	(± 7.8)	40.4	(± 7.8)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q8

Percent answers to 'Have you received a prescription for opioid pain medication from more than one healthcare provider?' by 'All Respondents'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 799)	10.6	(± 2.2)	89.1	(± 2.2)	0.3	(± 0.4)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you received a prescription for opioid pain medication from more than one healthcare provider?' by 'Age'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Age 18-35 (N = 263)	11.8	(± 4.0)	87.5	(± 4.1)	0.8	(± 1.1)
Age 36-64 (N = 376)	11.2	(± 3.3)	88.8	(± 3.3)	0.0	(± 0.0)
Age 65+ (N = 160)	7.5	(± 4.2)	92.5	(± 4.2)	0.0	(± 0.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you received a prescription for opioid pain medication from more than one healthcare provider?' by 'Gender'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Female (N = 403)	8.4	(± 2.8)	91.6	(± 2.8)	0.0	(± 0.0)
Male (N = 394)	12.9	(± 3.4)	86.5	(± 3.4)	0.5	(± 0.7)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you received a prescription for opioid pain medication from more than one healthcare provider?' by 'Race'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Black (N = 28)	7.1	(± 9.7)	92.9	(± 9.7)	0.0	(± 0.0)
Hispanic (N = 207)	9.2	(± 4.0)	90.8	(± 4.0)	0.0	(± 0.0)
Other race (N = 76)	10.5	(± 7.0)	89.5	(± 7.0)	0.0	(± 0.0)
White (N = 488)	11.5	(± 2.9)	88.1	(± 2.9)	0.4	(± 0.6)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_Q9

Percent answers to 'Have you ever used any prescription drugs without a doctor telling you to take them?' by 'All Respondents'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 799)	8.3	(± 1.9)	91.5	(± 2.0)	0.3	(± 0.4)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you ever used any prescription drugs without a doctor telling you to take them?' by 'Age'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Age 18-35 (N = 263)	11.8	(± 4.0)	87.8	(± 4.0)	0.4	(± 0.8)

Age 36-64 (N = 376)	7.7	(± 2.8)	92.0	(± 2.8)	0.3	(± 0.5)
Age 65+ (N = 160)	3.8	(± 3.0)	96.2	(± 3.0)	0.0	(± 0.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you ever used any prescription drugs without a doctor telling you to take them?' by 'Gender'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Female (N = 404)	6.9	(± 2.5)	93.1	(± 2.5)	0.0	(± 0.0)
Male (N = 393)	9.7	(± 3.0)	89.8	(± 3.1)	0.5	(± 0.7)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you ever used any prescription drugs without a doctor telling you to take them?' by 'Race'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Black (N = 28)	7.1	(± 9.7)	92.9	(± 9.7)	0.0	(± 0.0)
Hispanic (N = 208)	7.7	(± 3.7)	91.8	(± 3.8)	0.5	(± 1.0)
Other race (N = 76)	10.5	(± 7.0)	89.5	(± 7.0)	0.0	(± 0.0)
White (N = 487)	8.2	(± 2.5)	91.6	(± 2.5)	0.2	(± 0.4)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_Q10

Percent answers to 'Is it more difficult to get a pain medication prescription now than it was a few year ago?' by 'All Respondents'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 799)	34.9	(± 3.4)	27.5	(± 3.2)	37.5	(± 3.4)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Is it more difficult to get a pain medication prescription now than it was a few year ago?' by 'Age'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Age 18-35 (N = 264)	29.5	(± 5.6)	25.8	(± 5.4)	44.7	(± 6.1)
Age 36-64 (N = 375)	40.5	(± 5.1)	26.7	(± 4.6)	32.8	(± 4.9)
Age 65+ (N = 160)	30.6	(± 7.3)	32.5	(± 7.4)	36.9	(± 7.6)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Is it more difficult to get a pain medication prescription now than it was a few year ago?' by 'Gender'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Female (N = 404)	34.9	(± 4.7)	28.5	(± 4.5)	36.6	(± 4.8)
Male (N = 393)	34.9	(± 4.8)	26.7	(± 4.5)	38.4	(± 4.9)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Is it more difficult to get a pain medication prescription now than it was a few year ago?' by 'Race'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Black (N = 28)	25.0	(±16.4)	17.9	(±14.5)	57.1	(±18.7)
Hispanic (N = 208)	38.5	(± 6.8)	26.4	(± 6.1)	35.1	(± 6.6)
Other race (N = 76)	31.6	(±10.7)	22.4	(± 9.6)	46.1	(±11.4)
White (N = 487)	34.5	(± 4.3)	29.4	(± 4.1)	36.1	(± 4.4)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_Q11

Percent answers to 'Do you personally know anyone who has ever taken a prescription painkiller that was not prescribed to them, or not?' by 'All Respondents'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 800)	44.8	(± 3.5)	53.4	(± 3.5)	1.9	(± 1.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Do you personally know anyone who has ever taken a prescription painkiller that was not prescribed to them, or not?' by 'Age'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Age 18-35 (N = 264)	58.0	(± 6.1)	39.8	(± 6.0)	2.3	(± 1.8)
Age 36-64 (N = 376)	44.7	(± 5.1)	53.2	(± 5.1)	2.1	(± 1.5)
Age 65+ (N = 160)	23.1	(± 6.7)	76.2	(± 6.7)	0.6	(± 1.2)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Do you personally know anyone who has ever taken a prescription painkiller that was not prescribed to them, or not?' by 'Gender'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Female (N = 404)	41.8	(± 4.9)	55.9	(± 4.9)	2.2	(± 1.5)
Male (N = 394)	47.7	(± 5.0)	50.8	(± 5.0)	1.5	(± 1.2)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Do you personally know anyone who has ever taken a prescription painkiller that was not prescribed to them, or not?' by 'Race'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Black (N = 28)	42.9	(±18.7)	53.6	(±18.9)	3.6	(± 7.0)
Hispanic (N = 208)	38.0	(± 6.7)	60.6	(± 6.8)	1.4	(± 1.7)
Other race (N = 76)	53.9	(±11.4)	44.7	(±11.4)	1.3	(± 2.6)
White (N = 488)	46.3	(± 4.5)	51.6	(± 4.5)	2.0	(± 1.3)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q12_cata_1

Percent answers to 'Yourself--Who do you know that has ever taken a prescription painkiller that was not prescribed to them? Is it....

' by 'All Respondents'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
All Respondents (N = 343)	91.5	(± 3.0)	8.5	(± 3.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Yourself--Who do you know that has ever taken a prescription painkiller that was not prescribed to them? Is it....

' by 'Age'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Age 18-35 (N = 150)	90.7	(± 4.8)	9.3	(± 4.8)
Age 36-64 (N = 158)	91.8	(± 4.4)	8.2	(± 4.4)
Age 65+ (N = 35)	94.3	(± 7.9)	5.7	(± 7.9)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Yourself--Who do you know that has ever taken a prescription painkiller that was not prescribed to them? Is it....

' by 'Gender'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Female (N = 159)	92.5	(± 4.2)	7.5	(± 4.2)
Male (N = 183)	90.7	(± 4.3)	9.3	(± 4.3)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Yourself--Who do you know that has ever taken a prescription painkiller that was not prescribed to them? Is it....

' by 'Race'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Black (N = 11)	81.8	(±23.3)	18.2	(±23.3)
Hispanic (N = 76)	96.1	(± 4.5)	3.9	(± 4.5)
Other race (N = 40)	90.0	(± 9.5)	10.0	(± 9.5)
White (N = 216)	90.7	(± 4.0)	9.3	(± 4.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_12_cata_2

Percent answers to 'A family member--Who do you know that has ever taken a prescription painkiller that was not prescribed to them? Is it....

' by 'All Respondents'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
All Respondents (N = 343)	66.2	(± 5.1)	33.8	(± 5.1)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'A family member--Who do you know that has ever taken a prescription painkiller that was not prescribed to them? Is it....

' by 'Age'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Age 18-35 (N = 150)	64.0	(± 7.8)	36.0	(± 7.8)
Age 36-64 (N = 158)	67.1	(± 7.5)	32.9	(± 7.5)
Age 65+ (N = 35)	71.4	(±15.3)	28.6	(±15.3)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'A family member--Who do you know that has ever taken a prescription painkiller that was not prescribed to them? Is it....

' by 'Gender'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Female (N = 159)	59.7	(± 7.8)	40.3	(± 7.8)
Male (N = 183)	71.6	(± 6.7)	28.4	(± 6.7)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'A family member--Who do you know that has ever taken a prescription painkiller that was not prescribed to them? Is it....

' by 'Race'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Black (N = 11)	63.6	(±29.1)	36.4	(±29.1)
Hispanic (N = 76)	75.0	(± 9.9)	25.0	(± 9.9)
Other race (N = 40)	80.0	(±12.7)	20.0	(±12.7)
White (N = 216)	60.6	(± 6.7)	39.4	(± 6.7)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q12_cata_3

Percent answers to 'A close friend--Who do you know that has ever taken a prescription painkiller that was not prescribed to them? Is it....

' by 'All Respondents'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
All Respondents (N = 343)	58.6	(± 5.3)	41.4	(± 5.3)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'A close friend--Who do you know that has ever taken a prescription painkiller that was not prescribed to them? Is it....

' by 'Age'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Age 18-35 (N = 150)	54.0	(± 8.2)	46.0	(± 8.2)
Age 36-64 (N = 158)	60.8	(± 7.8)	39.2	(± 7.8)
Age 65+ (N = 35)	68.6	(±15.7)	31.4	(±15.7)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'A close friend--Who do you know that has ever taken a prescription painkiller that was not prescribed to them? Is it....

' by 'Gender'

Not Checked	Checked
-------------	---------

	Percent	MOE	Percent	MOE
Female (N = 159)	59.7	(± 7.8)	40.3	(± 7.8)
Male (N = 183)	57.4	(± 7.3)	42.6	(± 7.3)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'A close friend--Who do you know that has ever taken a prescription painkiller that was not prescribed to them? Is it....

' by 'Race'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Black (N = 11)	36.4	(±29.1)	63.6	(±29.1)
Hispanic (N = 76)	61.8	(±11.2)	38.2	(±11.2)
Other race (N = 40)	57.5	(±15.7)	42.5	(±15.7)
White (N = 216)	58.8	(± 6.7)	41.2	(± 6.7)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q12_cata_4

Percent answers to 'An acquaintance--Who do you know that has ever taken a prescription painkiller that was not prescribed to them? Is it....

' by 'All Respondents'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
All Respondents (N = 343)	48.4	(± 5.4)	51.6	(± 5.4)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'An acquaintance--Who do you know that has ever taken a prescription painkiller that was not prescribed to them? Is it....

' by 'Age'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Age 18-35 (N = 150)	44.0	(± 8.1)	56.0	(± 8.1)
Age 36-64 (N = 158)	51.3	(± 8.0)	48.7	(± 8.0)
Age 65+ (N = 35)	54.3	(±16.9)	45.7	(±16.9)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'An acquaintance--Who do you know that has ever taken a prescription painkiller that was not prescribed to them? Is it....

' by 'Gender'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE

Female (N = 159)	56.0	(± 7.9)	44.0	(± 7.9)
Male (N = 183)	42.1	(± 7.3)	57.9	(± 7.3)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'An acquaintance--Who do you know that has ever taken a prescription painkiller that was not prescribed to them? Is it....

' by 'Race'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Black (N = 11)	54.5	(±30.1)	45.5	(±30.1)
Hispanic (N = 76)	48.7	(±11.5)	51.3	(±11.5)
Other race (N = 40)	37.5	(±15.3)	62.5	(±15.3)
White (N = 216)	50.0	(± 6.8)	50.0	(± 6.8)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q13

Percent answers to 'When did you last use any prescription drugs without a doctor telling you to take them?										
' by 'All Respondents'										
			In the last 30 days		In the last 12 months but not within the last 30 days		Sometime in your lifetime but not within the last 12 months		(DON'T READ) Don't know / Not sure	
			Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 29)			3.4	(± 6.9)	27.6	(±16.9)	65.5	(±18.0)	3.4	(± 6.9)
Notes: 95 percent margin of error (MOE) in parentheses.										
Percent answers to 'When did you last use any prescription drugs without a doctor telling you to take them?'										
' by 'Age'										
			In the last 30 days		In the last 12 months but not within the last 30 days		Sometime in your lifetime but not within the last 12 months		(DON'T READ) Don't know / Not sure	
			Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE
Age 18-35 (N = 14)			0.0	(± 0.0)	28.6	(±24.6)	64.3	(±26.1)	7.1	(±14.0)
Age 36-64 (N = 13)			7.7	(±15.0)	30.8	(±26.1)	61.5	(±27.5)	0.0	(± 0.0)

Age 65+ (N = 2)	0.0	(± 0.0)	0.0	(± 0.0)	100.0	(± 0.0)	0.0	(± 0.0)	
Notes: 95 percent margin of error (MOE) in parentheses.									
Percent answers to 'When did you last use any prescription drugs without a doctor telling you to take them?' ' by 'Gender'									
	In the last 30 days		In the last 12 months but not within the last 30 days		Sometime in your lifetime but not within the last 12 months		(DON'T READ) Don't know / Not sure		
	Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE	
Female (N = 12)	8.3	(±16.2)	33.3	(±27.7)	58.3	(±29.0)	0.0	(± 0.0)	
Male (N = 17)	0.0	(± 0.0)	23.5	(±20.9)	70.6	(±22.5)	5.9	(±11.6)	
Notes: 95 percent margin of error (MOE) in parentheses.									
Percent answers to 'When did you last use any prescription drugs without a doctor telling you to take them?' ' by 'Race'									
	In the last 30 days		In the last 12 months but not within the last 30 days		Sometime in your lifetime but not within the last 12 months		(DON'T READ) Don't know / Not sure		
	Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE	
Black (N = 2)	0.0	(± 0.0)	0.0	(± 0.0)	100.0	(± 0.0)	0.0	(± 0.0)	
Hispanic (N = 3)	0.0	(± 0.0)	0.0	(± 0.0)	100.0	(± 0.0)	0.0	(± 0.0)	
Other race (N = 4)	0.0	(± 0.0)	25.0	(±44.1)	75.0	(±44.1)	0.0	(± 0.0)	
White (N = 20)	5.0	(± 9.9)	35.0	(±21.7)	55.0	(±22.6)	5.0	(± 9.9)	
Notes: 95 percent margin of error (MOE) in parentheses.									

Tab_q14

Percent answers to 'Do you personally know anyone who has ever been addicted to prescription painkillers, or not?' by 'All Respondents'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 799)	41.8	(± 3.5)	57.1	(± 3.5)	1.1	(± 0.7)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Do you personally know anyone who has ever been addicted to prescription painkillers, or not?' by 'Age'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Age 18-35 (N = 264)	45.1	(± 6.1)	54.2	(± 6.1)	0.8	(± 1.1)
Age 36-64 (N = 376)	43.4	(± 5.1)	55.1	(± 5.1)	1.6	(± 1.3)
Age 65+ (N = 159)	32.7	(± 7.4)	66.7	(± 7.5)	0.6	(± 1.3)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Do you personally know anyone who has ever been addicted to prescription painkillers, or not?' by 'Gender'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Female (N = 404)	39.9	(± 4.9)	58.9	(± 4.9)	1.2	(± 1.1)
Male (N = 393)	43.8	(± 5.0)	55.2	(± 5.0)	1.0	(± 1.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Do you personally know anyone who has ever been addicted to prescription painkillers, or not?' by 'Race'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Black (N = 28)	17.9	(±14.5)	78.6	(±15.5)	3.6	(± 7.0)
Hispanic (N = 208)	33.2	(± 6.5)	66.3	(± 6.6)	0.5	(± 1.0)
Other race (N = 76)	44.7	(±11.4)	55.3	(±11.4)	0.0	(± 0.0)
White (N = 487)	46.4	(± 4.5)	52.2	(± 4.5)	1.4	(± 1.1)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q15_cata_1

Percent answers to 'Yourself--Who do you know that has ever been addicted to prescription painkillers? Is it...

' by 'All Respondents'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
All Respondents (N = 325)	95.4	(± 2.3)	4.6	(± 2.3)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Yourself--Who do you know that has ever been addicted to prescription painkillers? Is it...

' by 'Age'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Age 18-35 (N = 118)	99.2	(± 1.7)	0.8	(± 1.7)
Age 36-64 (N = 158)	93.7	(± 3.9)	6.3	(± 3.9)
Age 65+ (N = 49)	91.8	(± 7.8)	8.2	(± 7.8)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Yourself--Who do you know that has ever been addicted to prescription painkillers? Is it...

' by 'Gender'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Female (N = 154)	97.4	(± 2.6)	2.6	(± 2.6)
Male (N = 170)	93.5	(± 3.8)	6.5	(± 3.8)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Yourself--Who do you know that has ever been addicted to prescription painkillers? Is it...

' by 'Race'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Black (N = 5)	100.0	(± 0.0)	0.0	(± 0.0)
Hispanic (N = 68)	94.1	(± 5.7)	5.9	(± 5.7)
Other race (N = 34)	97.1	(± 5.8)	2.9	(± 5.8)
White (N = 218)	95.4	(± 2.8)	4.6	(± 2.8)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q5_cata_2

Percent answers to 'A family member--Who do you know that has ever been addicted to prescription painkillers? Is it...

' by 'All Respondents'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
All Respondents (N = 325)	57.8	(± 5.5)	42.2	(± 5.5)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'A family member--Who do you know that has ever been addicted to prescription painkillers? Is it...

' by 'Age'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Age 18-35 (N = 118)	57.6	(± 9.1)	42.4	(± 9.1)
Age 36-64 (N = 158)	60.8	(± 7.8)	39.2	(± 7.8)
Age 65+ (N = 49)	49.0	(±14.3)	51.0	(±14.3)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'A family member--Who do you know that has ever been addicted to prescription painkillers? Is it...

' by 'Gender'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Female (N = 154)	53.9	(± 8.0)	46.1	(± 8.0)
Male (N = 170)	61.2	(± 7.5)	38.8	(± 7.5)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'A family member--Who do you know that has ever been addicted to prescription painkillers? Is it...

' by 'Race'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Black (N = 5)	100.0	(± 0.0)	0.0	(± 0.0)
Hispanic (N = 68)	69.1	(±11.2)	30.9	(±11.2)
Other race (N = 34)	61.8	(±16.7)	38.2	(±16.7)
White (N = 218)	52.8	(± 6.8)	47.2	(± 6.8)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q15_cata_3

Percent answers to 'A close friend--Who do you know that has ever been addicted to prescription painkillers? Is it...

' by 'All Respondents'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
All Respondents (N = 325)	69.2	(± 5.1)	30.8	(± 5.1)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'A close friend--Who do you know that has ever been addicted to prescription painkillers? Is it...

' by 'Age'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Age 18-35 (N = 118)	67.8	(± 8.6)	32.2	(± 8.6)
Age 36-64 (N = 158)	67.7	(± 7.5)	32.3	(± 7.5)
Age 65+ (N = 49)	77.6	(±11.9)	22.4	(±11.9)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'A close friend--Who do you know that has ever been addicted to prescription painkillers? Is it...

' by 'Gender'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Female (N = 154)	70.1	(± 7.4)	29.9	(± 7.4)
Male (N = 170)	68.2	(± 7.2)	31.8	(± 7.2)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'A close friend--Who do you know that has ever been addicted to prescription painkillers? Is it...

' by 'Race'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Black (N = 5)	20.0	(±35.8)	80.0	(±35.8)
Hispanic (N = 68)	72.1	(±10.9)	27.9	(±10.9)
Other race (N = 34)	70.6	(±15.7)	29.4	(±15.7)
White (N = 218)	69.3	(± 6.3)	30.7	(± 6.3)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q15_cata_4

Percent answers to 'An acquaintance--Who do you know that has ever been addicted to prescription painkillers? Is it...

' by 'All Respondents'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
All Respondents (N = 325)	54.5	(± 5.5)	45.5	(± 5.5)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'An acquaintance--Who do you know that has ever been addicted to prescription painkillers? Is it...

' by 'Age'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Age 18-35 (N = 118)	54.2	(± 9.2)	45.8	(± 9.2)
Age 36-64 (N = 158)	53.2	(± 8.0)	46.8	(± 8.0)
Age 65+ (N = 49)	59.2	(±14.1)	40.8	(±14.1)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'An acquaintance--Who do you know that has ever been addicted to prescription painkillers? Is it...

' by 'Gender'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Female (N = 154)	56.5	(± 8.0)	43.5	(± 8.0)
Male (N = 170)	52.9	(± 7.7)	47.1	(± 7.7)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'An acquaintance--Who do you know that has ever been addicted to prescription painkillers? Is it...

' by 'Race'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Black (N = 5)	60.0	(±43.9)	40.0	(±43.9)
Hispanic (N = 68)	54.4	(±12.1)	45.6	(±12.1)
Other race (N = 34)	50.0	(±17.2)	50.0	(±17.2)
White (N = 218)	55.0	(± 6.7)	45.0	(± 6.7)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q16

Percent answers to 'Do you personally know anyone who has died from a prescription painkiller overdose, or not?' by 'All Respondents'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 799)	14.4	(± 2.5)	85.4	(± 2.5)	0.3	(± 0.4)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Do you personally know anyone who has died from a prescription painkiller overdose, or not?' by 'Age'

Yes No Don't Know

	Percent	MOE	Percent	MOE	Percent	MOE
Age 18-35 (N = 264)	15.5	(± 4.5)	83.7	(± 4.5)	0.8	(± 1.1)
Age 36-64 (N = 376)	15.4	(± 3.7)	84.6	(± 3.7)	0.0	(± 0.0)
Age 65+ (N = 159)	10.1	(± 4.8)	89.9	(± 4.8)	0.0	(± 0.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Do you personally know anyone who has died from a prescription painkiller overdose, or not?' by 'Gender'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Female (N = 404)	13.9	(± 3.4)	86.1	(± 3.4)	0.0	(± 0.0)
Male (N = 393)	15.0	(± 3.6)	84.5	(± 3.7)	0.5	(± 0.7)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Do you personally know anyone who has died from a prescription painkiller overdose, or not?' by 'Race'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Black (N = 28)	3.6	(± 7.0)	96.4	(± 7.0)	0.0	(± 0.0)
Hispanic (N = 208)	13.5	(± 4.7)	86.5	(± 4.7)	0.0	(± 0.0)
Other race (N = 76)	14.5	(± 8.1)	85.5	(± 8.1)	0.0	(± 0.0)
White (N = 487)	15.4	(± 3.3)	84.2	(± 3.3)	0.4	(± 0.6)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q17_cata_1

Percent answers to 'A family member--Who do you know that has died from a prescription painkiller overdose? Is it...

' by 'All Respondents'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
All Respondents (N = 113)	74.3	(± 8.3)	25.7	(± 8.3)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'A family member--Who do you know that has died from a prescription painkiller overdose? Is it...

' by 'Age'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Age 18-35 (N = 40)	85.0	(±11.3)	15.0	(±11.3)
Age 36-64 (N = 58)	69.0	(±12.2)	31.0	(±12.2)
Age 65+ (N = 15)	66.7	(±24.5)	33.3	(±24.5)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'A family member--Who do you know that has died from a prescription painkiller overdose? Is it...

' by 'Gender'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Female (N = 55)	70.9	(±12.3)	29.1	(±12.3)
Male (N = 58)	77.6	(±11.0)	22.4	(±11.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'A family member--Who do you know that has died from a prescription painkiller overdose? Is it...

' by 'Race'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Black (N = 1)	100.0	(± 0.0)	0.0	(± 0.0)
Hispanic (N = 28)	71.4	(±17.2)	28.6	(±17.2)
Other race (N = 11)	90.9	(±17.4)	9.1	(±17.4)
White (N = 73)	72.6	(±10.5)	27.4	(±10.5)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q17_cata_2

Percent answers to 'A close friend--Who do you know that has died from a prescription painkiller overdose? Is it...

' by 'All Respondents'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
All Respondents (N = 113)	68.1	(± 8.8)	31.9	(± 8.8)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'A close friend--Who do you know that has died from a prescription painkiller overdose? Is it...

' by 'Age'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Age 18-35 (N = 40)	77.5	(±13.3)	22.5	(±13.3)
Age 36-64 (N = 58)	63.8	(±12.7)	36.2	(±12.7)
Age 65+ (N = 15)	60.0	(±25.4)	40.0	(±25.4)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'A close friend--Who do you know that has died from a prescription painkiller overdose? Is it...

' by 'Gender'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Female (N = 55)	72.7	(±12.1)	27.3	(±12.1)
Male (N = 58)	63.8	(±12.7)	36.2	(±12.7)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'A close friend--Who do you know that has died from a prescription painkiller overdose? Is it...

' by 'Race'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Black (N = 1)	100.0	(± 0.0)	0.0	(± 0.0)
Hispanic (N = 28)	85.7	(±13.3)	14.3	(±13.3)
Other race (N = 11)	45.5	(±30.2)	54.5	(±30.2)
White (N = 73)	64.4	(±11.3)	35.6	(±11.3)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q17_cata_3

Percent answers to 'An acquaintance--Who do you know that has died from a prescription painkiller overdose? Is it...

' by 'All Respondents'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
All Respondents (N = 113)	50.4	(± 9.4)	49.6	(± 9.4)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'An acquaintance--Who do you know that has died from a prescription painkiller overdose? Is it...

' by 'Age'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Age 18-35 (N = 40)	32.5	(±14.9)	67.5	(±14.9)
Age 36-64 (N = 58)	58.6	(±13.0)	41.4	(±13.0)
Age 65+ (N = 15)	66.7	(±24.5)	33.3	(±24.5)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'An acquaintance--Who do you know that has died from a prescription painkiller overdose? Is it...

' by 'Gender'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Female (N = 55)	50.9	(±13.5)	49.1	(±13.5)
Male (N = 58)	50.0	(±13.2)	50.0	(±13.2)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'An acquaintance--Who do you know that has died from a prescription painkiller overdose? Is it...

' by 'Race'

	Not Checked		Checked	
	Percent	MOE	Percent	MOE
Black (N = 1)	0.0	(± 0.0)	100.0	(± 0.0)
Hispanic (N = 28)	42.9	(±18.8)	57.1	(±18.8)
Other race (N = 11)	45.5	(±30.2)	54.5	(±30.2)
White (N = 73)	54.8	(±11.7)	45.2	(±11.7)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q18

Percent answers to 'In the past 12 months, have you, personally, spent more time drinking or using drugs than you intended?' by 'All Respondents'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 798)	3.6	(± 1.3)	96.1	(± 1.4)	0.3	(± 0.4)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'In the past 12 months, have you, personally, spent more time drinking or using drugs than you intended?' by 'Age'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Age 18-35 (N = 264)	3.0	(± 2.1)	97.0	(± 2.1)	0.0	(± 0.0)
Age 36-64 (N = 376)	4.8	(± 2.2)	94.9	(± 2.3)	0.3	(± 0.5)
Age 65+ (N = 158)	1.9	(± 2.2)	97.5	(± 2.5)	0.6	(± 1.3)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'In the past 12 months, have you, personally, spent more time drinking or using drugs than you intended?' by 'Gender'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Female (N = 404)	2.2	(± 1.5)	97.8	(± 1.5)	0.0	(± 0.0)
Male (N = 392)	5.1	(± 2.2)	94.4	(± 2.3)	0.5	(± 0.7)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'In the past 12 months, have you, personally, spent more time drinking or using drugs than you intended?' by 'Race'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Black (N = 28)	0.0	(± 0.0)	100.0	(± 0.0)	0.0	(± 0.0)
Hispanic (N = 208)	5.3	(± 3.1)	94.7	(± 3.1)	0.0	(± 0.0)
Other race (N = 76)	2.6	(± 3.7)	97.4	(± 3.7)	0.0	(± 0.0)
White (N = 486)	3.3	(± 1.6)	96.3	(± 1.7)	0.4	(± 0.6)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q19

Percent answers to 'How many times have you been prescribed a prescription pain reliever in the past 12 months?' by 'All Respondents'												
			Once		Two or three times		Four or more times		Never		Don't Know	
			Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 798)			17.8	(± 2.7)	8.1	(± 1.9)	5.8	(± 1.7)	67.4	(± 3.3)	0.9	(± 0.7)
Notes: 95 percent margin of error (MOE) in parentheses.												
Percent answers to 'How many times have you been prescribed a prescription pain reliever in the past 12 months?' by 'Age'												
			Once		Two or three times		Four or more times		Never		Don't Know	
			Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE
Age 18-35 (N = 263)			17.5	(± 4.7)	6.8	(± 3.1)	1.1	(± 1.3)	73.8	(± 5.4)	0.8	(± 1.1)
Age 36-64 (N = 375)			18.1	(± 4.0)	10.7	(± 3.2)	8.0	(± 2.8)	61.9	(± 5.0)	1.3	(± 1.2)

Age 65+ (N = 160)	17.5	(± 6.0)	4.4	(± 3.2)	8.1	(± 4.3)	70.0	(± 7.3)	0.0	(± 0.0)	
Notes: 95 percent margin of error (MOE) in parentheses.											
Percent answers to 'How many times have you been prescribed a prescription pain reliever in the past 12 months?' by 'Gender'											
	Once		Two or three times		Four or more times		Never		Don't Know		
	Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE	
Female (N = 403)	18.4	(± 3.9)	9.2	(± 2.9)	6.0	(± 2.4)	65.0	(± 4.8)	1.5	(± 1.2)	
Male (N = 393)	17.3	(± 3.8)	7.1	(± 2.6)	5.3	(± 2.3)	70.0	(± 4.6)	0.3	(± 0.5)	
Notes: 95 percent margin of error (MOE) in parentheses.											
Percent answers to 'How many times have you been prescribed a prescription pain reliever in the past 12 months?' by 'Race'											
	Once		Two or three times		Four or more times		Never		Don't Know		
	Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE	
Black (N = 28)	7.1	(± 9.7)	10.7	(±11.7)	7.1	(± 9.7)	71.4	(±17.1)	3.6	(± 7.0)	
Hispanic (N = 207)	19.8	(± 5.5)	5.3	(± 3.1)	2.4	(± 2.1)	70.0	(± 6.4)	2.4	(± 2.1)	
Other race (N = 76)	17.1	(± 8.6)	5.3	(± 5.1)	5.3	(± 5.1)	72.4	(±10.3)	0.0	(± 0.0)	
White (N = 487)	17.7	(± 3.5)	9.7	(± 2.7)	7.2	(± 2.3)	65.3	(± 4.3)	0.2	(± 0.4)	
Notes: 95 percent margin of error (MOE) in parentheses.											

Tab_q20

Percent answers to 'Approximately how many times in the past 12 months have you combined alcohol and sedatives like valium or Xanax?' by 'All Respondents'											
	Once		Two or three times		Four or more times		Never		Don't Know		
	Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE	
All Respond	0.6	(± 0.6)	0.9	(± 0.7)	0.1	(± 0.3)	98.1	(± 1.0)	0.3	(± 0.4)	

ents (N = 797)											
Notes: 95 percent margin of error (MOE) in parentheses.											
Percent answers to 'Approximately how many times in the past 12 months have you combined alcohol and sedatives like valium or Xanax?' by 'Age'											
		Once		Two or three times		Four or more times		Never		Don't Know	
		Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE
		t	(±)	t	(±)	t	(±)	t	(±)	t	(±)
Age 18-35 (N = 262)		0.8	(± 1.1)	1.5	(± 1.5)	0.0	(± 0.0)	97.7	(± 1.8)	0.0	(± 0.0)
Age 36-64 (N = 376)		0.8	(± 0.9)	0.3	(± 0.5)	0.3	(± 0.5)	98.4	(± 1.3)	0.3	(± 0.5)
Age 65+ (N = 159)		0.0	(± 0.0)	1.3	(± 1.8)	0.0	(± 0.0)	98.1	(± 2.2)	0.6	(± 1.3)
Notes: 95 percent margin of error (MOE) in parentheses.											
Percent answers to 'Approximately how many times in the past 12 months have you combined alcohol and sedatives like valium or Xanax?' by 'Gender'											
		Once		Two or three times		Four or more times		Never		Don't Know	
		Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE
		t	(±)	t	(±)	t	(±)	t	(±)	t	(±)
Female (N = 403)		0.2	(± 0.5)	0.2	(± 0.5)	0.0	(± 0.0)	99.3	(± 0.9)	0.2	(± 0.5)
Male (N = 392)		1.0	(± 1.0)	1.5	(± 1.2)	0.3	(± 0.5)	96.9	(± 1.7)	0.3	(± 0.5)
Notes: 95 percent margin of error (MOE) in parentheses.											
Percent answers to 'Approximately how many times in the past 12 months have you combined alcohol and sedatives like valium or Xanax?' by 'Race'											
		Once		Two or three times		Four or more times		Never		Don't Know	
		Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE
		t	(±)	t	(±)	t	(±)	t	(±)	t	(±)
Black (N = 28)		0.0	(± 0.0)	0.0	(± 0.0)	0.0	(± 0.0)	100.0	(± 0.0)	0.0	(± 0.0)
Hispanic (N = 206)		1.9	(± 1.9)	0.5	(± 1.0)	0.0	(± 0.0)	97.6	(± 2.1)	0.0	(± 0.0)

Other race (N = 76)	0.0	(± 0.0)	1.3	(± 2.6)	1.3	(± 2.6)	97.4	(± 3.7)	0.0	(± 0.0)
White (N = 487)	0.2	(± 0.4)	1.0	(± 0.9)	0.0	(± 0.0)	98.4	(± 1.2)	0.4	(± 0.6)
Notes: 95 percent margin of error (MOE) in parentheses.										

Tab_q21

Percent answers to 'Have you ever not been able to perform normal functions as a result of misusing a prescription pain reliever?' by 'All Respondents'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 796)	4.6	(± 1.5)	94.8	(± 1.6)	0.5	(± 0.5)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you ever not been able to perform normal functions as a result of misusing a prescription pain reliever?' by 'Age'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Age 18-35 (N = 262)	4.6	(± 2.6)	95.4	(± 2.6)	0.0	(± 0.0)
Age 36-64 (N = 375)	5.6	(± 2.4)	93.9	(± 2.5)	0.5	(± 0.8)
Age 65+ (N = 159)	2.5	(± 2.5)	96.2	(± 3.0)	1.3	(± 1.8)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you ever not been able to perform normal functions as a result of misusing a prescription pain reliever?' by 'Gender'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Female (N = 402)	3.2	(± 1.8)	96.3	(± 1.9)	0.5	(± 0.7)
Male (N = 392)	5.9	(± 2.4)	93.6	(± 2.5)	0.5	(± 0.7)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you ever not been able to perform normal functions as a result of misusing a prescription pain reliever?' by 'Race'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Black (N = 27)	7.4	(±10.1)	92.6	(±10.1)	0.0	(± 0.0)
Hispanic (N = 207)	3.9	(± 2.7)	95.7	(± 2.8)	0.5	(± 1.0)
Other race (N = 76)	5.3	(± 5.1)	94.7	(± 5.1)	0.0	(± 0.0)

White (N = 486)	4.7	(± 1.9)	94.7	(± 2.0)	0.6	(± 0.7)
-----------------	-----	---------	------	---------	-----	---------

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q22

Percent answers to 'Has anyone ever told you they were worried about your misuse of prescription drugs?' by 'All Respondents'

	Yes		No	
	Percent	MOE	Percent	MOE
All Respondents (N = 799)	3.0	(± 1.2)	97.0	(± 1.2)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Has anyone ever told you they were worried about your misuse of prescription drugs?' by 'Age'

	Yes		No	
	Percent	MOE	Percent	MOE
Age 18-35 (N = 264)	1.5	(± 1.5)	98.5	(± 1.5)
Age 36-64 (N = 375)	3.5	(± 1.9)	96.5	(± 1.9)
Age 65+ (N = 160)	4.4	(± 3.2)	95.6	(± 3.2)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Has anyone ever told you they were worried about your misuse of prescription drugs?' by 'Gender'

	Yes		No	
	Percent	MOE	Percent	MOE
Female (N = 403)	1.7	(± 1.3)	98.3	(± 1.3)
Male (N = 394)	4.3	(± 2.0)	95.7	(± 2.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Has anyone ever told you they were worried about your misuse of prescription drugs?' by 'Race'

	Yes		No	
	Percent	MOE	Percent	MOE
Black (N = 27)	0.0	(± 0.0)	100.0	(± 0.0)
Hispanic (N = 208)	3.4	(± 2.5)	96.6	(± 2.5)
Other race (N = 76)	0.0	(± 0.0)	100.0	(± 0.0)
White (N = 488)	3.5	(± 1.7)	96.5	(± 1.7)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q23

Percent answers to 'Has your doctor ever talked to you about the risk or dependence upon prescription drugs?' by 'All Respondents'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 800)	27.9	(± 3.2)	71.4	(± 3.2)	0.8	(± 0.6)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Has your doctor ever talked to you about the risk or dependence upon prescription drugs?' by 'Age'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Age 18-35 (N = 264)	27.3	(± 5.5)	71.6	(± 5.6)	1.1	(± 1.3)
Age 36-64 (N = 376)	32.7	(± 4.8)	66.8	(± 4.9)	0.5	(± 0.8)
Age 65+ (N = 160)	17.5	(± 6.0)	81.9	(± 6.1)	0.6	(± 1.2)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Has your doctor ever talked to you about the risk or dependence upon prescription drugs?' by 'Gender'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Female (N = 404)	24.3	(± 4.3)	75.5	(± 4.3)	0.2	(± 0.5)
Male (N = 394)	31.2	(± 4.7)	67.5	(± 4.7)	1.3	(± 1.1)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Has your doctor ever talked to you about the risk or dependence upon prescription drugs?' by 'Race'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Black (N = 28)	35.7	(±18.1)	64.3	(±18.1)	0.0	(± 0.0)
Hispanic (N = 208)	32.2	(± 6.5)	66.3	(± 6.6)	1.4	(± 1.7)
Other race (N = 76)	23.7	(± 9.8)	76.3	(± 9.8)	0.0	(± 0.0)
White (N = 488)	26.2	(± 4.0)	73.2	(± 4.0)	0.6	(± 0.7)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q24

Percent answers to 'Have you ever been screened for depression or anxiety?' by 'All Respondents'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 800)	28.0	(± 3.2)	71.8	(± 3.2)	0.2	(± 0.4)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you ever been screened for depression or anxiety?' by 'Age'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Age 18-35 (N = 264)	26.9	(± 5.5)	72.7	(± 5.5)	0.4	(± 0.8)
Age 36-64 (N = 376)	32.7	(± 4.8)	67.0	(± 4.9)	0.3	(± 0.5)
Age 65+ (N = 160)	18.8	(± 6.2)	81.2	(± 6.2)	0.0	(± 0.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you ever been screened for depression or anxiety?' by 'Gender'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Female (N = 404)	31.4	(± 4.6)	68.3	(± 4.6)	0.2	(± 0.5)
Male (N = 394)	24.4	(± 4.3)	75.4	(± 4.3)	0.3	(± 0.5)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you ever been screened for depression or anxiety?' by 'Race'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Black (N = 28)	28.6	(±17.1)	71.4	(±17.1)	0.0	(± 0.0)
Hispanic (N = 208)	20.7	(± 5.6)	78.8	(± 5.7)	0.5	(± 1.0)
Other race (N = 76)	30.3	(±10.5)	69.7	(±10.5)	0.0	(± 0.0)
White (N = 488)	30.7	(± 4.2)	69.1	(± 4.2)	0.2	(± 0.4)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q25

Percent answers to 'Do you lock up your medication?' by 'All Respondents'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 793)	25.3	(± 3.1)	72.8	(± 3.2)	1.9	(± 1.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Do you lock up your medication?' by 'Age'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Age 18-35 (N = 262)	27.9	(± 5.5)	70.2	(± 5.7)	1.9	(± 1.7)
Age 36-64 (N = 374)	30.2	(± 4.8)	68.4	(± 4.8)	1.3	(± 1.2)
Age 65+ (N = 157)	9.6	(± 4.7)	87.3	(± 5.3)	3.2	(± 2.8)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Do you lock up your medication?' by 'Gender'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Female (N = 401)	26.9	(± 4.4)	71.3	(± 4.5)	1.7	(± 1.3)
Male (N = 390)	23.8	(± 4.3)	74.1	(± 4.4)	2.1	(± 1.4)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Do you lock up your medication?' by 'Race'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Black (N = 27)	22.2	(±16.0)	74.1	(±16.9)	3.7	(± 7.3)
Hispanic (N = 206)	36.4	(± 6.7)	62.6	(± 6.7)	1.0	(± 1.4)
Other race (N = 76)	17.1	(± 8.6)	80.3	(± 9.1)	2.6	(± 3.7)
White (N = 484)	22.1	(± 3.8)	75.8	(± 3.9)	2.1	(± 1.3)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q26

Percent answers to 'Do you know how to properly store pain medication?' by 'All Respondents'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 797)	65.2	(± 3.4)	29.5	(± 3.2)	5.3	(± 1.6)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Do you know how to properly store pain medication?' by 'Age'

Yes No Don't Know

	Percent	MOE	Percent	MOE	Percent	MOE
Age 18-35 (N = 264)	61.0	(± 6.0)	35.6	(± 5.9)	3.4	(± 2.2)
Age 36-64 (N = 375)	71.2	(± 4.7)	24.3	(± 4.4)	4.5	(± 2.1)
Age 65+ (N = 158)	58.2	(± 7.9)	31.6	(± 7.4)	10.1	(± 4.8)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Do you know how to properly store pain medication?' by 'Gender'

	Yes Percent	MOE	No Percent	MOE	Don't Know Percent	MOE
Female (N = 404)	66.6	(± 4.7)	25.7	(± 4.4)	7.7	(± 2.7)
Male (N = 391)	63.7	(± 4.9)	33.5	(± 4.8)	2.8	(± 1.7)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Do you know how to properly store pain medication?' by 'Race'

	Yes Percent	MOE	No Percent	MOE	Don't Know Percent	MOE
Black (N = 28)	64.3	(±18.1)	25.0	(±16.4)	10.7	(±11.7)
Hispanic (N = 208)	60.1	(± 6.8)	35.6	(± 6.6)	4.3	(± 2.8)
Other race (N = 76)	61.8	(±11.2)	36.8	(±11.1)	1.3	(± 2.6)
White (N = 485)	68.0	(± 4.2)	26.0	(± 4.0)	6.0	(± 2.2)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q27

Percent answers to 'Have you ever worried about overdosing?' by 'All Respondents'

	Yes Percent	MOE	No Percent	MOE	Don't Know Percent	MOE
All Respondents (N = 799)	6.8	(± 1.8)	93.1	(± 1.8)	0.1	(± 0.3)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you ever worried about overdosing?' by 'Age'

	Yes Percent	MOE	No Percent	MOE	Don't Know Percent	MOE
Age 18-35 (N = 264)	7.6	(± 3.3)	92.4	(± 3.3)	0.0	(± 0.0)

Age 36-64 (N = 376)	8.0	(± 2.8)	91.8	(± 2.8)	0.3	(± 0.5)
Age 65+ (N = 159)	2.5	(± 2.5)	97.5	(± 2.5)	0.0	(± 0.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you ever worried about overdosing?' by 'Gender'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Female (N = 404)	5.7	(± 2.3)	94.1	(± 2.4)	0.2	(± 0.5)
Male (N = 393)	7.9	(± 2.7)	92.1	(± 2.7)	0.0	(± 0.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you ever worried about overdosing?' by 'Race'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Black (N = 28)	0.0	(± 0.0)	96.4	(± 7.0)	3.6	(± 7.0)
Hispanic (N = 208)	14.4	(± 4.9)	85.6	(± 4.9)	0.0	(± 0.0)
Other race (N = 76)	3.9	(± 4.5)	96.1	(± 4.5)	0.0	(± 0.0)
White (N = 487)	4.3	(± 1.8)	95.7	(± 1.8)	0.0	(± 0.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q28

Percent answers to 'Have you ever overdosed?' by 'All Respondents'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 798)	2.4	(± 1.1)	97.5	(± 1.1)	0.1	(± 0.3)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you ever overdosed?' by 'Age'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Age 18-35 (N = 263)	0.8	(± 1.1)	99.2	(± 1.1)	0.0	(± 0.0)
Age 36-64 (N = 376)	4.0	(± 2.0)	95.7	(± 2.1)	0.3	(± 0.5)

Age 65+ (N = 159)	1.3	(± 1.8)	98.7	(± 1.8)	0.0	(± 0.0)
-------------------	-----	---------	------	---------	-----	---------

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you ever overdosed?' by 'Gender'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Female (N = 403)	1.7	(± 1.3)	98.0	(± 1.4)	0.2	(± 0.5)
Male (N = 393)	3.1	(± 1.7)	96.9	(± 1.7)	0.0	(± 0.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you ever overdosed?' by 'Race'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Black (N = 28)	0.0	(± 0.0)	96.4	(± 7.0)	3.6	(± 7.0)
Hispanic (N = 207)	3.9	(± 2.7)	96.1	(± 2.7)	0.0	(± 0.0)
Other race (N = 76)	2.6	(± 3.7)	97.4	(± 3.7)	0.0	(± 0.0)
White (N = 487)	1.8	(± 1.2)	98.2	(± 1.2)	0.0	(± 0.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q29

Percent answers to 'Have you ever used heroin?' by 'All Respondents'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 795)	1.9	(± 1.0)	98.0	(± 1.0)	0.1	(± 0.3)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you ever used heroin?' by 'Age'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Age 18-35 (N = 263)	0.4	(± 0.8)	99.6	(± 0.8)	0.0	(± 0.0)
Age 36-64 (N = 374)	3.5	(± 1.9)	96.3	(± 2.0)	0.3	(± 0.5)
Age 65+ (N = 158)	0.6	(± 1.3)	99.4	(± 1.3)	0.0	(± 0.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you ever used heroin?' by 'Gender'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Female (N = 401)	0.7	(± 0.9)	99.0	(± 1.0)	0.2	(± 0.5)
Male (N = 392)	2.8	(± 1.7)	97.2	(± 1.7)	0.0	(± 0.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you ever used heroin?' by 'Race'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Black (N = 28)	0.0	(± 0.0)	96.4	(± 7.0)	3.6	(± 7.0)
Hispanic (N = 206)	1.9	(± 1.9)	98.1	(± 1.9)	0.0	(± 0.0)
Other race (N = 76)	3.9	(± 4.5)	96.1	(± 4.5)	0.0	(± 0.0)
White (N = 485)	1.6	(± 1.2)	98.4	(± 1.2)	0.0	(± 0.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q30

Percent answers to 'Are you, yourself, now covered by any form of health insurance or health plan or do you not have health insurance at this time?

' by 'All Respondents'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 793)	92.4	(± 1.9)	7.1	(± 1.8)	0.5	(± 0.5)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Are you, yourself, now covered by any form of health insurance or health plan or do you not have health insurance at this time?

' by 'Age'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Age 18-35 (N = 263)	90.1	(± 3.7)	9.5	(± 3.6)	0.4	(± 0.8)
Age 36-64 (N = 372)	92.5	(± 2.7)	6.7	(± 2.6)	0.8	(± 0.9)
Age 65+ (N = 158)	96.2	(± 3.0)	3.8	(± 3.0)	0.0	(± 0.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Are you, yourself, now covered by any form of health insurance or health plan or do you not have health insurance at this time?

' by 'Gender'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Female (N = 402)	94.0	(± 2.4)	5.5	(± 2.3)	0.5	(± 0.7)
Male (N = 389)	91.0	(± 2.9)	8.5	(± 2.8)	0.5	(± 0.7)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Are you, yourself, now covered by any form of health insurance or health plan or do you not have health insurance at this time?

' by 'Race'

	Yes		No		Don't Know	
	Percent	MOE	Percent	MOE	Percent	MOE
Black (N = 28)	78.6	(±15.5)	14.3	(±13.2)	7.1	(± 9.7)
Hispanic (N = 206)	88.8	(± 4.4)	10.2	(± 4.2)	1.0	(± 1.4)
Other race (N = 76)	88.2	(± 7.4)	11.8	(± 7.4)	0.0	(± 0.0)
White (N = 483)	95.4	(± 1.9)	4.6	(± 1.9)	0.0	(± 0.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q31

Percent answers to 'Which of the following is your main source of health insurance coverage?																	
' by 'All Respondents'																	
		A plan through your employer		A plan through your spouse's employer		A plan you purchased yourself		Medicare		Medicaid		A plan through a parent or parents		Or from some other source?		(DON'T READ) Don't Know	
		Per cent	MOE	Per cent	MOE	Per cent	MOE	Per cent	MOE	Per cent	MOE	Per cent	MOE	Per cent	MOE	Per cent	MOE
All Respondents		37.0	(± 3.6)	7.4	(± 1.9)	7.9	(± 2.0)	24.8	(± 3.2)	6.4	(± 1.8)	3.9	(± 1.4)	11.5	(± 2.4)	1.1	(± 0.8)

(N = 721)																		
Notes: 95 percent margin of error (MOE) in parentheses.																		
Percent answers to 'Which of the following is your main source of health insurance coverage?'																		
' by 'Age'																		
			A plan through your employer		A plan through your spouse's employer		A plan you purchased yourself		Medicare		Medicaid		A plan through a parent or parents		Or from some other source?		(DON'T READ) Don't Know	
			Per cent	MOE	Per cent	MOE	Per cent	MOE	Per cent	MOE	Per cent	MOE	Per cent	MOE	Per cent	MOE	Per cent	MOE
		Age 18-35 (N = 234)	52.1	(± 6.5)	11.1	(± 4.1)	7.3	(± 3.4)	3.8	(± 2.5)	6.0	(± 3.1)	10.7	(± 4.0)	8.1	(± 3.6)	0.9	(± 1.2)
		Age 36-64 (N = 340)	40.0	(± 5.3)	6.8	(± 2.7)	9.1	(± 3.1)	19.4	(± 4.3)	8.5	(± 3.0)	0.9	(± 1.0)	15.0	(± 3.9)	0.3	(± 0.6)
		Age 65+ (N = 147)	6.1	(± 4.0)	2.7	(± 2.7)	6.1	(± 4.0)	70.7	(± 7.5)	2.0	(± 2.3)	0.0	(± 0.0)	8.8	(± 4.7)	3.4	(± 3.0)
Notes: 95 percent margin of error (MOE) in parentheses.																		
Percent answers to 'Which of the following is your main source of health insurance coverage?'																		
' by 'Gender'																		
			A plan through your employer		A plan through your spouse's employer		A plan you purchased yourself		Medicare		Medicaid		A plan through a parent or parents		Or from some other source?		(DON'T READ) Don't Know	
			Per cent	MOE	Per cent	MOE	Per cent	MOE	Per cent	MOE	Per cent	MOE	Per cent	MOE	Per cent	MOE	Per cent	MOE
		Female (N = 371)	35.6	(± 5.0)	10.0	(± 3.1)	6.7	(± 2.6)	27.2	(± 4.6)	7.5	(± 2.7)	3.0	(± 1.8)	8.9	(± 3.0)	1.1	(± 1.1)

Male (N = 349)	38.7	(± 5.2)	4.6	(± 2.2)	9.2	(± 3.1)	22.3	(± 4.5)	4.9	(± 2.3)	4.9	(± 2.3)	14.3	(± 3.8)	1.1	(± 1.1)
Notes: 95 percent margin of error (MOE) in parentheses.																
Percent answers to 'Which of the following is your main source of health insurance coverage?' ' by 'Race'																
			A plan through your employer	A plan through your spouse's employer	A plan you purchased yourself	Medicare	Medicaid	A plan through a parent or parents	Or from some other source?	(DON'T READ) Don't Know						
			Per cent	MOE	Per cent	MOE	Per cent	MOE	Per cent	MOE	Per cent	MOE	Per cent	MOE	Per cent	MOE
Black (N = 22)	31.8	(± 9.9)	4.5	(± 8.9)	13.6	(± 4.6)	13.6	(± 4.6)	4.5	(± 8.9)	31.8	(± 19.9)	0.0	(± 0.0)	0.0	(± 0.0)
Hispanic (N = 182)	41.8	(± 7.3)	7.7	(± 4.0)	5.5	(± 3.4)	14.3	(± 5.2)	9.9	(± 4.4)	4.4	(± 3.0)	14.8	(± 5.3)	1.6	(± 1.9)
Other race (N = 67)	44.8	(± 2.2)	19.4	(± 9.7)	4.5	(± 5.1)	13.4	(± 8.3)	4.5	(± 5.1)	4.5	(± 5.1)	9.0	(± 7.0)	0.0	(± 0.0)
White (N = 450)	34.2	(± 4.5)	5.6	(± 2.2)	9.1	(± 2.7)	31.3	(± 4.4)	5.3	(± 2.1)	2.2	(± 1.4)	11.1	(± 3.0)	1.1	(± 1.0)
Notes: 95 percent margin of error (MOE) in parentheses.																

Tab_q32

Percent answers to 'Have you been pregnant in the past 12 months?' by 'All Respondents'

	Yes	MOE	No	MOE
	Percent		Percent	
All Respondents (N = 173)	13.3	(± 5.2)	86.7	(± 5.2)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you been pregnant in the past 12 months?' by 'Age'

	Yes Percent	MOE	No Percent	MOE
Age 18-35 (N = 127)	17.3	(± 6.7)	82.7	(± 6.7)
Age 36-64 (N = 46)	2.2	(± 4.3)	97.8	(± 4.3)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you been pregnant in the past 12 months?' by 'Gender'

	Yes Percent	MOE	No Percent	MOE
Female (N = 173)	13.3	(± 5.2)	86.7	(± 5.2)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Have you been pregnant in the past 12 months?' by 'Race'

	Yes Percent	MOE	No Percent	MOE
Black (N = 11)	0.0	(± 0.0)	100.0	(± 0.0)
Hispanic (N = 60)	8.3	(± 7.2)	91.7	(± 7.2)
Other race (N = 20)	15.0	(±16.0)	85.0	(±16.0)
White (N = 82)	18.3	(± 8.6)	81.7	(± 8.6)

Notes: 95 percent margin of error (MOE) in parentheses.

tab_q33

Percent answers to 'Did you take any prescription pain relievers during pregnancy?' by 'All Respondents'

	No Percent	MOE
All Respondents (N = 23)	100.0	(± 0.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Did you take any prescription pain relievers during pregnancy?' by 'Age'

	No Percent	MOE
Age 18-35 (N = 22)	100.0	(± 0.0)
Age 36-64 (N = 1)	100.0	(± 0.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Did you take any prescription pain relievers during pregnancy?' by 'Gender'

	No	MOE
	Percent	(± 0.0)
Female (N = 23)	100.0	(± 0.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Did you take any prescription pain relievers during pregnancy?' by 'Race'

	No	MOE
	Percent	(± 0.0)
Hispanic (N = 5)	100.0	(± 0.0)
Other race (N = 3)	100.0	(± 0.0)
White (N = 15)	100.0	(± 0.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_q34

Percent answers to 'Did your doctor talk to you about possible consequences of taking prescription pain relievers during pregnancy?' by 'All Respondents'

	Yes	MOE	No	MOE
	Percent	(±21.3)	Percent	(±21.3)
All Respondents (N = 23)	47.8	(±21.3)	52.2	(±21.3)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Did your doctor talk to you about possible consequences of taking prescription pain relievers during pregnancy?' by 'Age'

	Yes	MOE	No	MOE
	Percent	(±21.8)	Percent	(±21.8)
Age 18-35 (N = 22)	50.0	(±21.8)	50.0	(±21.8)
Age 36-64 (N = 1)	0.0	(± 0.0)	100.0	(± 0.0)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Did your doctor talk to you about possible consequences of taking prescription pain relievers during pregnancy?' by 'Gender'

	Yes	MOE	No	MOE
	Percent	(±21.3)	Percent	(±21.3)
Female (N = 23)	47.8	(±21.3)	52.2	(±21.3)

Notes: 95 percent margin of error (MOE) in parentheses.

Percent answers to 'Did your doctor talk to you about possible consequences of taking prescription pain relievers during pregnancy?' by 'Race'

	Yes	MOE	No	MOE
	Percent	(±44.8)	Percent	(±44.8)
Hispanic (N = 5)	60.0	(±44.8)	40.0	(±44.8)
Other race (N = 3)	33.3	(±55.7)	66.7	(±55.7)
White (N = 15)	46.7	(±26.3)	53.3	(±26.3)

Notes: 95 percent margin of error (MOE) in parentheses.

Tab_race

Percent answers to 'Race' by 'All Respondents'									
Black				Hispanic		Other race		White	
		Percent	MOE	Percent	MOE	Percent	MOE	Percent	MOE
All Respondents (N = 800)		3.5	(± 1.3)	26.0	(± 3.1)	9.5	(± 2.1)	61.0	(± 3.5)
Notes: 95 percent margin of error (MOE) in parentheses.									

Tab_sex

Percent answers to 'Gender' by 'All Respondents'

	Female		Male	
	Percent	MOE	Percent	MOE
All Respondents (N = 798)	50.6	(± 3.5)	49.4	(± 3.5)

Notes: 95 percent margin of error (MOE) in parentheses.