



TOXICOLOGY LABORATORY
WASHINGTON STATE PATROL
2203 Airport Way South Suite 360 Seattle, WA 98134
(206) 262-6100 FAX No. (206) 262-6145

JUL 31 REC'D

TOXICOLOGY TEST REPORT

MB 7137

Attention: Dr. Brian Mazrim
Agency: King Co Medical Examiner
Address: Box 359792
325 9th Ave
Seattle, WA 98104-2499

Tox Case #: ST-17-07273 Case Type: Death Investigation Report Date: 7/26/2017

Agency Case #: 17-01206 Subject Name: TENT: Charleena C. Lyles

Evidence: The following evidence was submitted to the Laboratory by Zachary Porras-Foye of the King Co Medical Examiner on 6/20/2017 via hand delivery:
(1) ST-17-07273-A: VGreen, Blood - Hospital

Volatile Analysis Results:

ST-17-07273-A: Blood - Hospital

ST-17-07273-A was tested by Headspace - Gas Chromatography for the presence of acetone, ethanol, isopropanol, and methanol on 07/13/2017. The following result was obtained:

None Detected

Drug Analysis Results:

ST-17-07273-A: Blood - Hospital

ST-17-07273-A was tested by Enzyme Multiplied Immunoassay Technique (EMIT) for the presence of amphetamines, barbiturates, benzodiazepines, cannabinoids, cocaine metabolite, and opiates on 07/14/2017. The following result(s) was obtained:

None Detected


ST-17-07273-A was tested by Gas Chromatography/Mass Spectrometry for the presence of basic drugs and metabolites on 07/20/2017. The following result(s) was obtained:

None Detected

COMMENTS

All testing was performed by the Forensic Scientist listed below except as otherwise indicated. The Forensic Scientist has technically reviewed all relevant pages of testing documentation in the case record.

CONFIDENTIAL
DO NOT COPY
R.C.W. 06.00.105





Case Number: 17-1206

Name: _____

Date: 6-19-17

Pathologist: BM

A) Gunshot exit wound:
0.5x0.5" defect ±
0.2x0.2" abrasion ±
inf/med margin.
3.25" @ mid,
49" sup to heel.

Grazing gunshot
wound:
2.5x1" defect
~ lat @ chest
13" sup to heel.
inf skin tags post/post
Gunshot entrance
wound: 0.4"
circ defect ±
conc. 0.05"
abrasion margin.
5" @ mid,
0" sup to heel.
No suture/stitching.

Gunshot entrance
wound: 0.35"
circ defect ±
0.05 to 0.2" abrasion
margin, accent sup.
entered midline,
0.5" sup to heel.

Gunshot exit
wound: 0.6" diam.
irreg defect on
ant axillary line,
12.5" sup to heel.


Gunshot entrance wound:
0.35" circ defect ±
conc. 0.05" abrasion margin.
Ext @ hip, 34.5" sup to heel.
No suture/stitching.

G) Gunshot entrance wound:
0.3" circ defect ±
conc. 0.1" abrasion margin.
lateral @ arm 45"
sup to heel. No suture/stitching.

H) Gunshot exit wound:
0.8x0.6" stellate defect
on postero-medial
@ arm, 45" sup
to heel.

I) Gunshot entrance wound:
0.25" circ defect ±
conc. 0.1" abrasion
margin. Centered 0.5"
@ mid, 46.25" sup to heel.
No suture/stitching.

J) Gunshot entrance
wound: 0.5x0.3"
defect ± eccentric
abrasion margin,
eccent up to
0.5", accent
medially. Centered
3.75" @ mid,
42" sup to
heel. No suture/
stitching.



CONFIDENTIAL
DO NOT COPY
R.C.W. 68.50.105